

Port of Brisbane Weed Surveys 2018

Reference: R.B20259.027.01.2018 weed survey.docx
Date: June 2018
Confidential



Document Control Sheet

BMT WBM Pty Ltd Level 8, 200 Creek Street Brisbane Qld 4000 Australia PO Box 203, Spring Hill 4004 Tel: +61 7 3831 6744 Fax: + 61 7 3832 3627 ABN 54 010 830 421 www.bmt.org	Document:	R.B20259.027.01.2018 weed survey.docx
	Title:	Port of Brisbane Weed Surveys 2018
	Project Manager:	Suanne Richards
	Author:	Suanne Richards and Sophie Hipkin
	Client:	Port of Brisbane
	Client Contact:	Jessica Rudd
	Client Reference:	
Synopsis: This report presents the findings of the Port of Brisbane Annual Weed Monitoring Program for 2018.		

REVISION/CHECKING HISTORY

Revision Number	Date	Checked by	Issued by
0	15 June 2018	DLR	SR
1	27 June 2018	DLR	SR

DISTRIBUTION

Destination	Revision										
	0	1	2	3	4	5	6	7	8	9	10
PBPL	PDF	PDF									
BMT File	PDF	PDF									
BMT Library	PDF	PDF									

Copyright and non-disclosure notice

The contents and layout of this report are subject to copyright owned by BMT WBM Pty Ltd (BMT WBM) save to the extent that copyright has been legally assigned by us to another party or is used by BMT WBM under licence. To the extent that we own the copyright in this report, it may not be copied or used without our prior written agreement for any purpose other than the purpose indicated in this report.

The methodology (if any) contained in this report is provided to you in confidence and must not be disclosed or copied to third parties without the prior written agreement of BMT WBM. Disclosure of that information may constitute an actionable breach of confidence or may otherwise prejudice our commercial interests. Any third party who obtains access to this report by any means will, in any event, be subject to the Third Party Disclaimer set out below.

Third Party Disclaimer

Any disclosure of this report to a third party is subject to this disclaimer. The report was prepared by BMT WBM at the instruction of, and for use by, our client named on this Document Control Sheet. It does not in any way constitute advice to any third party who is able to access it by any means. BMT WBM excludes to the fullest extent lawfully permitted all liability whatsoever for any loss or damage howsoever arising from reliance on the contents of this report.

Executive Summary

Executive Summary

The Port of Brisbane Annual Weed Monitoring Program has been conducted on Port of Brisbane Pty Ltd (PBPL) lands since 2001. The monitoring program aims to identify the introduction and spread of priority weed species on PBPL lands and to recommend management and control measures as required.

This report provides the results for the 2018 weed monitoring program for survey sites at Lucinda Drain, Port Gate Drain, Port West Drain, T1-3, Car Precinct and the constructed lake, and sites of high environmental value and/or at high risk of weed invasion.

All weed species recorded on PBPL lands are well-established in the Brisbane region and south-east Queensland. No targeted National priority pest plants listed by the Department of Agriculture and Water Resources (DAWR) were recorded within or directly adjacent to the survey areas.

The most abundant weed species recorded on PBPL lands included:


- Woody weeds: Broad leaf peppertree *Schinus terebinthifolius*, which was common on disturbed terrestrial lands, and has also invaded saltmarsh communities. *Leucaena leucocephala* has also become more widespread in some saltmarsh habitats.
- Vine cover: Siratro *Macroptilium atropurpureum* was the most widespread vine species recorded on PBPL lands. This species had variable abundance where it occurred, and in places formed dense patches (50-90% cover) resulting in smothering of native vegetation.
- Ground cover: The highest weed species richness was recorded in the ground-layer, reflecting the higher richness of groundcover weed species in the region. Highest weed groundcover (50-90%) was generally recorded in disturbed terrestrial sites. The most abundant groundcover weed species

were green panic *Megathyrsus maximus* var. *maximus*, Rhode's grass *Chloris gayana* and red natal grass *Melinis repens*.

Weed invasion on PBPL lands generally occurs within and directly adjacent to disturbed lands, within terrestrial lands and at the limit of Highest Astronomical Tide (HAT). Habitats subject to regular tidal inundation are unaffected by weed invasion. The marine weed mangrove species *Rhizophora mangle* has been recorded in Australia but has not been observed in south-east Queensland and is therefore considered a low risk species. Despite the low risk, regular weed monitoring should be continued to ensure this species, and other high-risk taxa, do not establish in Port habitats.

Survey sites have remained relatively stable over the monitoring period (2001-2018). Except for isolated thatch grass *Hyparrhenia rufa* (a widespread grass weed in south-east Qld), no new weed species, or significant weed invasion outbreaks, were recorded in the 2018 survey. The weed species recorded on PBPL lands are widespread in Brisbane and the south-east Queensland bioregion, and for that reason will be difficult to control and are unlikely to be eradicated from PBPL lands.

It is recommended that PBPL continue weed monitoring and control efforts to fulfil landholder obligations under Queensland's *Biosecurity Act*. This includes monitoring the Port Drive corridor and sites of high ecological value, especially saltmarsh, which is listed as a vulnerable ecological community under Commonwealth legislation. Saltmarsh is sensitive to a range of threats including land disturbance, hydrological modifications, weed invasion and mangrove encroachment resulting from sea level rise. Improving land management practices east of Port Drive and re-profiling exotic grasslands in this area would also assist saltmarsh resilience to ongoing threats. Ongoing weed monitoring and management will also reduce potential future costs associated with delaying weed control, particularly the removal of mature woody shrubs and dense vine infestations.



Site Summary

- Bird Hide:**

 - intertidal saltmarsh providing important habitat for migratory waders
 - some Phragmites encroachment into saltmarsh
 - Restricted Matters under Biosecurity Act: groundsel
- Lucinda Drain:**

 - constructed intertidal channel with maintained banks
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree, camphor laurel, lantanas, prickly pear
- T1-3, Car Precinct and Constructed Lakes:**

 - constructed lakes and regularly mowed easements
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: groundsel, fireweed, ragweed, broad-leaved pepper tree, lantana
- Port Drive (North) Wetlands:**


 - mangroves and saltmarsh associated with Boat Passage
 - saltmarsh-mangrove ecotone supports localised patches of broad-leaved pepper tree and exotic grasses
 - landward edge of saltmarsh is regularly slashed, reducing habitat condition and extent of this threatened community
 - Phragmites encroachment into saltmarsh
 - re-profiling highly disturbed areas of exotic grassland adjacent to tidal lands could assist saltmarsh resilience to sea level rise and may promote recruitment of nationally threatened swamp oak communities
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree
- Port Gate Drain:**

 - constructed intertidal channel
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: groundsel, broad-leaved pepper tree, lantana
- Fort Lytton Estuarine Wetlands:**

 - mangroves and saltmarsh associated with the Brisbane River
 - saltmarsh-mangrove ecotone supports localised patches of broad-leaved pepper tree and exotic grasses
 - Phragmites encroachment into saltmarsh
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree, Chinese elm, groundsel, lantana, balloon vine, prickly pear, madeira vine
- Port West Estuarine Wetlands:**

 - mangroves and saltmarsh associated with the Brisbane River
 - saltmarsh-mangrove ecotone supports localised patches of broad-leaved pepper tree and exotic grasses
 - Phragmites encroachment into saltmarsh
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree, ground asparagus fern, prickly pear
- Port West Drain:**


 - constructed intertidal channel
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: groundsel, prickly pear, broad-leaved pepper tree, lantana, madeira vine



- LEGEND**
- Weed Monitoring Sites**
- | | |
|--------------------------------|--|
| <div></div> Bird Hide | <div></div> Port Gate Drain |
| <div></div> Fort Lytton | <div></div> Port West |
| <div></div> Lucinda Drain | <div></div> Port West Drain |
| <div></div> Port Drive - North | <div></div> T1-3 / Car Precinct / Lake |

Title:
2018 Weed Monitoring Sites

BMT WBM endeavours to ensure that the information provided in this map is correct at the time of publication. BMT WBM does not warrant, guarantee or make representations regarding the currency and accuracy of information contained in this map.



Filepath: I:\B20259_I_BRH Port of Brisbane DLR\DRG\Weeds 2018\DRG\ECO_004_2018 Weed Summary Descriptive.WOR

Figure:
1-1

Rev:
A



Contents**Contents**

Executive Summary	i
1 Introduction	1
1.1 Background	1
1.2 Aims	2
2 Methodology	4
2.1 Targeted Weed Species	4
2.2 Techniques	5
2.3 Survey Sites	5
2.4 Survey Limitations	5
3 Results	7
3.1 Lucinda Drain	7
3.1.1 Site Description	7
3.1.2 Weeds	7
3.2 Port West Drain	11
3.2.1 Site Description	11
3.2.2 Weeds	11
3.3 Port Gate Drain	12
3.3.1 Site Description	12
3.3.2 Weeds	12
3.4 T1-3, Car Precinct and Constructed Lakes	14
3.4.1 Site Description	14
3.4.2 Weeds	15
3.5 The Bird Hide	16
3.6 Port Drive Wetlands	17
3.6.1 Intertidal Wetlands	17
3.6.2 Filled Intertidal Land	18
3.6.3 Freshwater Wetlands	20
3.7 Port West Estuarine Wetlands	20
3.8 Fort Lytton Estuarine Wetlands	21
4 Discussion	23
5 Conclusion	25
6 References	26
Appendix A Dominant Weed Species Recorded at PBPL	A-1

Contents**List of Figures**

Figure 1-1	2018 Weed Monitoring Sites	3
Figure 3-1	Lucinda Drain 2018	8
Figure 3-2	2018 Weed Survey Summary	9
Figure 3-3	Port West Drain 2018	12
Figure 3-4	Port Gate Drain 2018	14
Figure 3-5	T1-3, Car Precinct and Constructed Lake	16
Figure 3-6	Bird Hide 2018	17
Figure 3-7	Disturbed Coastal Saltmarsh Port Drive	19
Figure 3-8	Port Drive Wetlands 2018	20
Figure 3-9	Port West Estuarine Habitat 2018	21
Figure 3-10	Fort Lytton 2018	22
Figure 4-1	Mangrove Encroachment of Coastal Saltmarsh Port Drive 2009-2018	24

1 Introduction

1.1 Background

Annual weed monitoring surveys have been conducted at the Port of Brisbane (PBPL) since 2001. The weed monitoring program aims to detect and assess the potential introduction and spread of priority weed species on PBPL lands and to recommend management and control measures as required. Priority weeds targeted in the surveys include plant species considered to be exotic or invasive, namely:

- Weeds listed by the Commonwealth Department of Agriculture and Water Resources (DAWR). DAWR listed weed species are specifically targeted because of their potential to enter the country via containers and other materials shipped and unloaded at the PBPL.
- Prohibited and Restricted Matters regulated under the Queensland *Biosecurity Act 2014*.
- Environmental weeds listed by the Brisbane City Council (BCC).

Surveys have been conducted at Lucinda Drain from 2001-2018, with surveys at Port Gate Drain commencing in 2007. In 2013 Port West was included in the survey area due to the identification of pest species by stakeholders. In response to a potential weed incursion threat from imported vehicles, weed survey sites at the Car Precinct and T1-3 Overflow Area were included in the monitoring program in 2008. However, given regular maintenance of the Car Precinct and Overflow Areas the risk of weed invasion and spread is considered low in this area and monitoring was reduced to biennial assessments from 2014. Additional sites of high environmental value and/or at high risk of weed invasion were included in the program in 2016 (BMT WBM 2016).

Based on previous survey results, weed composition and distribution at the Port have remained relatively stable over the monitoring period (BMT WBM 2017). All weeds recorded in the survey sites are widespread in the Brisbane region. No DAWR listed species have been recorded and although approximately 27 new plant species arrive in south east Queensland each year, no new weed species to the region have been recorded on Port lands. Restricted Matters recorded included: broad-leaf pepper tree (*Schinus terebinthifolius*), camphor laurel (*Cinnamomum camphora*), Chinese elm (*Celtis sinensis*), fireweed (*Senecio madagascariensis*), groundsel (*Baccharis halimifolia*), lantana (*Lantana camara*), creeping lantana (*Lantana montevidensis*), mother-of-millions (*Bryophyllum delagoense*), prickly pear (*Opuntia stricta*), giant paramatta grass (*Sporobolus fertilis*), annual ragweed (*Ambrosia artemisiifolia*), asparagus ferns (*Asparagus aethiopicus* and *A. africanus*) and balloon vine (*Cardiospermum grandiflorum*).

Several widespread environmental weeds listed by BCC have also been recorded in the survey sites. All weed species recorded at PBPL are widespread in degraded sites and coastal habitats of south-east Queensland. Refer to Appendix A for a list of dominant weeds recorded.

Introduction


1.2 Aims

Targeted weed surveys were conducted in May 2018 at Lucinda Drain, Port Gate Drain, Port West Drain, T1-3, Car Precinct, the constructed lake, and sites of high environmental value and/or at high risk of weed invasion identified by the BMT WBM (2016) (refer to Figure 1-1).

The freshwater wetlands adjacent to Port Drive were not surveyed in 2017 or 2018 due to access constraints associated with road widening and construction works.

The aims of the PBPL Weed Monitoring Program are to:

- Record new species or high risk weed infestations at monitoring sites; and
- Provide recommendations for strategic weed management at the Port.



Title:
2018 Weed Monitoring Sites

Figure:

1-1

Rev:

A

BMT WBM endeavours to ensure that the information provided in this map is correct at the time of publication. BMT WBM does not warrant, guarantee or make representations regarding the currency and accuracy of information contained in this map.



0km 0.8 1.6
Approx. Scale



2 Methodology

In accordance with previous monitoring surveys, weed inspections were undertaken in the post-summer months in May (2018). The survey was conducted by a qualified botanist with over 20 years pest survey experience in coastal south-east Queensland habitats.

2.1 Targeted Weed Species

Species targeted in the survey included the following:

- Weeds listed by DAWR (i.e National Environmental Alert List / Weeds of National Significance (WONS)), with the potential to invade south-east Queensland, including:
 - Chinese violet (*Asystasia gangetica*).
<http://www.environment.gov.au/biodiversity/invasive/weeds/publications/guidelines/alert/a-gangetica.html>
 - Siam weed (*Chromolaena odorata*).
<https://www.environment.gov.au/biodiversity/invasive/weeds/publications/guidelines/alert/pubs/c-odorata.pdf>
 - Cyperus sedge (*Cyperus teneristolon*).
http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=67172
 - Horsetails (*Equisetum hyemale*).
<https://www.environment.gov.au/biodiversity/invasive/weeds/publications/guidelines/alert/pubs/equisetum.pdf>
 - Senegal Tea Plant (*Gymnocoronis spilanthoides*).
http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=12850
 - Lagarosiphon (*Lagarosiphon major*).
<http://www.environment.gov.au/biodiversity/invasive/weeds/publications/guidelines/alert/pubs/l-major.pdf>
 - Leaf Cactus (*Pereskia aculeata*).
http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=12443
 - Praxelis (*Praxelis clematidea*).
http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=67811
 - Laurel Clock Vine (*Thunbergia laurifolia*).

Methodology

http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=42844

- Karroo Thorn (*Vachellia karroo*).

http://www.environment.gov.au/cgi-bin/biodiversity/invasive/weeds/weeddetails.pl?taxon_id=84350

- Prohibited and Restricted Matters regulated under the Queensland *Biosecurity Act 2014* - refer https://www.daf.qld.gov.au/_data/assets/pdf_file/0004/383818/IPA-Restricted-plants-of-Qld.pdf
- Species listed by BCC as environmental pests.
- 'High risk' species assessed by Biosecurity Queensland (DAF), including the potentially invasive mangrove *Rhizophora mangle*.

2.2 Techniques

The weed survey used the random meander technique for recording all exotic and invasive species within the survey sites. All surveys were conducted on-foot to ensure that extensive coverage of the survey sites was achieved. Incidental observations of targeted weed species outside the survey sites were also recorded. The locations of all observed target weed species were recorded on handheld GPS. Weed identification was undertaken on site. Samples of weed species unable to be identified *in situ* were pressed for later verification.

2.3 Survey Sites

Survey sites are indicated in Figure 1-1.

The weed survey along Lucinda Drain consisted of recording exotic and invasive species within a 15 m wide belt transect along the entire length of the drain's eastern bank. A visual inspection of the western bank was taken from the eastern bank.

The weed survey along Port West Drain consisted of recording exotic and invasive species within a 15 m wide belt transect along the entire length of the drain's eastern bank.

The weed survey along the Port Gate Drain consisted of recording exotic and invasive species within a 15 m wide transect along the entire length of the drain's eastern and western banks.

The weed survey in the T1-3 Overflow area consisted of recording exotic and invasive species within a 2 m wide transect along the entire eastern, northern, western and southern boundaries of the hardstand area and along both banks of the constructed drain just north of the hardstand area.

The weed survey of the Car Precinct Area consisted of recording exotic and invasive species within the easement between the railway and the western boundary of the hardstand area and the banks of the lake adjacent to the previous Visitors Centre site.

All sites of environmental significance were inspected on foot using the random meander technique.

2.4 Survey Limitations

Whilst every effort has been made to identify targeted weed species in the PBPL survey sites, the detectability of plant species and the ability to accurately identify these in the field varies with

Methodology

seasonal and climatic conditions. Such conditions influence the presence of reproductive features (flowers, fruits and seeds) which are useful, and in some cases essential, for species identification. Consequently, the survey conducted should not be regarded as conclusive that targeted weeds do not occur within the survey sites or surrounding lands.

3 Results

The following provides descriptions of weed composition and distribution recorded at the survey sites. Appendix A provides a summary of dominant weed species recorded on PBPL lands. Figure 3-2 summarises the findings from the 2018 weed survey. Only exotic or introduced flora species recorded during the surveys are detailed in the following site descriptions, unless otherwise noted.

3.1 Lucinda Drain

3.1.1 Site Description

Lucinda Drain is a constructed channel located east of Lucinda Drive on the eastern edge of the Port. It provides drainage for stormwater run-off from hardstand areas to the north and discharges through the Lucinda Weir into Boat Passage (refer to Figure 3-1).

The channel banks supported planted and naturally recruited shrubs and trees comprised of a mix of local native terrestrial species such as she-oaks (*Casuarina* spp.), figs (*Ficus* spp.), cotton tree (*Hibiscus tiliaceus*), *Melaleuca* spp. and *Macaranga tanarius*. Introduced shrubs were sparse but widespread and the groundcover was dominated by exotic grasses. The tidal channel does not contain extensive aquatic macrophytic cover but supported a low, discontinuous fringe of the native grey mangrove (*Avicennia marina*). The drain is periodically maintained and mangroves are actively removed to ensure the drain fulfils its primary purpose for stormwater run-off. Extensive mangroves and saltpan lie to the east of the drain associated with the intertidal flats of Boat Passage.

The western bank of the drain adjacent to Lucinda Drive undergoes regular maintenance involving mowing and weed spraying. Poor access along the eastern bank of Lucinda Drain limits regular maintenance but weeds are reportedly removed on an annual basis (RPS, 2013).

3.1.2 Weeds

The following observations were made in the 2018 survey:


- Lucinda Drain remains in a stable condition in terms of weed invasion.
- No DAWR-listed species were recorded within or directly adjacent to Lucinda Drain.
- Restricted Matters recorded included: broad-leaved pepper tree (*Schinus terebinthifolius*), camphor laurel (*Cinnamomum camphora*), lantana (*Lantana camara*), creeping lantana (*Lantana montevidensis*), prickly pear (*Opuntia stricta*) and giant rat's tail grass (*Sporobolus* sp.) were widespread but sparse along both banks of the channel.
- Exotic species recorded were dominated by environmental weeds well established across Brisbane including: *Opuntia stricta*, *Schinus terebinthifolius*, *Cinnamomum camphora*, *Lantana montevidensis*, Easter cassia (*Senna pendula* var. *glabrata*), siratro (*Macroptilium atropurpureum*), Rhode's grass (*Chloris gayana*), green panic (*Megathyrsus maximus* var. *maximus*), Brazilian nightshade (*Solanum seaforthianum*) and blackberry nightshade (*Solanum nigrum*).

Results

- Consistent with previous surveys, the dominant woody weed recorded at Lucinda Drain was *Schinus terebinthifolius*. Other woody weeds included *Lantana camara*, *Senna pendula* var. *glabrata* and *Leucaena leucocephala*.
- *Macroptilium atropurpureum*, mile a minute (*Ipomoea cairica*), glycine (*Neonotonia wightii*) and *Solanum seaforthianum* were the most abundant exotic vine species.
- The groundcover was dominated by mown exotic grasses including *Chloris gayana*, *Megathyrsus maximus* var. *maximus*, *Melinis repens* and Mossman River grass (*Cenchrus echinatus*).
- Other exotic groundcovers, forbs and herbs included *Stylosanthes scabra*, *Commelina benghalensis*, *Tridax procumbens*, *Calyptocarpus vialis*, *Gomphrena celosioides*, *Cynodon dactylon*, *Sida cordifolia*, *Capsella bursa-pastoris*, *Crotalaria pallida*, *Conyza bonariensis*, *Crassocephalum crepidioides*, *Bidens pilosa*, *Portulaca oleracea*, *Ageratum houstonianum*, *Oenothera drummondii*, *Plantago major*, *Sonchus oleraceus*, *Phyllanthus virgatus* and *Malvastrum coromandelianum*.
- No aquatic macrophyte weed species were recorded. The brackish to saline conditions of the channel limits the establishment of exotic aquatic macrophytes known from the region.
- No significant differences in weed composition or density have been observed for Lucinda Drain over time.



Figure 3-1 Lucinda Drain 2018



Site Summary

- Bird Hide:**
 - intertidal saltmarsh providing important habitat for migratory waders
 - some Phragmites encroachment into saltmarsh
 - Restricted Matters under Biosecurity Act: groundsel
- Lucinda Drain:**
 - constructed intertidal channel with maintained banks
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree, camphor laurel, lantanas, prickly pear
- T1-3, Car Precinct and Constructed Lakes:**
 - constructed lakes and regularly mowed easements
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: groundsel, fireweed, ragweed, broad-leaved pepper tree, lantana
- Port Drive (North) Wetlands:**
 - mangroves and saltmarsh associated with Boat Passage
 - saltmarsh-mangrove ecotone supports localised patches of broad-leaved pepper tree and exotic grasses
 - landward edge of saltmarsh is regularly slashed, reducing habitat condition and extent of this threatened community
 - Phragmites encroachment into saltmarsh
 - re-profiling highly disturbed areas of exotic grassland adjacent to tidal lands could assist saltmarsh resilience to sea level rise and may promote recruitment of nationally threatened swamp oak communities
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree
- Port Gate Drain:**
 - constructed intertidal channel
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: groundsel, broad-leaved pepper tree, lantana
- Fort Lytton Estuarine Wetlands:**
 - mangroves and saltmarsh associated with the Brisbane River
 - saltmarsh-mangrove ecotone supports localised patches of broad-leaved pepper tree and exotic grasses
 - Phragmites encroachment into saltmarsh
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree, Chinese elm, groundsel, lantana, balloon vine, prickly pear, madeira vine
- Port West Estuarine Wetlands:**
 - mangroves and saltmarsh associated with the Brisbane River
 - saltmarsh-mangrove ecotone supports localised patches of broad-leaved pepper tree and exotic grasses
 - Phragmites encroachment into saltmarsh
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: broad-leaved pepper tree, ground asparagus fern, prickly pear
- Port West Drain:**
 - constructed intertidal channel
 - stable condition
 - exotic species dominated by environmental weeds well established across Brisbane
 - Restricted Matters under Biosecurity Act: groundsel, prickly pear, broad-leaved pepper tree, lantana, madeira vine



LEGEND

Weed Monitoring Sites

- Bird Hide
- Fort Lytton
- Lucinda Drain
- Port Drive - North
- Port Gate Drain
- Port West
- Port West Drain
- T1-3 / Car Precinct / Lake

Title:

2018 Weed Monitoring Summary

BMT WBM endeavours to ensure that the information provided in this map is correct at the time of publication. BMT WBM does not warrant, guarantee or make representations regarding the currency and accuracy of information contained in this map.



0km 0.8 1.6
Approx. Scale

Figure:

3-2

Rev:

A



Filepath: I:\B20259 | BRH Port of Brisbane DLR\DRG\Weeds 2018\DRG\ECO_004_2018 Weed Summary Descriptive.WOR

3.2 Port West Drain

3.2.1 Site Description

Port West Drain, located west of Lytton Road approximately 4 km south-west of the Port, was comprised of a narrow intertidal channel fringed with remnant mangroves numerically dominated by mature *Avicennia marina* (refer to Figure 3-3). The channel is bounded to the west by extensive mangrove forest and cleared land for industrial purposes lie to the east. No scheduled maintenance work, including weed removal, is undertaken at Port West Drain (RPS, 2013).

3.2.2 Weeds

The following observations were made in the 2018 survey:

- No DAWR-listed species were recorded within or directly adjacent to Port West Drain.
- Weed species listed as Restricted Matters that were recorded here included: *Baccharis halimifolia*, *Opuntia stricta*, *Schinus terebinthifolius*, *Lantana camara* and madeira vine (*Anredera cordifolia*).
- Exotic species recorded included *Ipomoea cairica*, *Macroptilium atropurpureum*, *Chloris gayana*, *Megathyrsus maximus* var. *maximus*, *Bidens pilosa*, *Gomphocarpus physocarpus*, *Melinis repens*, *Solanum nigrum*, *Conyza bonariensis*, *Macroptilium lathyroides*, *Anredera cordifolia*, *Asparagus aethiopicus* and *Asparagus africanus*. These environmental weeds are well established across Brisbane.
- The dominant woody weeds recorded included *Schinus terebinthifolius*, *Baccharis halimifolia* and *Lantana camara*. Other woody weeds included *Gomphocarpus physocarpus*, phasey bean (*Macroptilium lathyroides*), castor oil plant (*Ricinus communis*), coral berry (*Rivina humilis*), *Senna pendula* var. *glabrata*, giant Devil's fig (*Solanum chrysotrichum*), wild tobacco (*Solanum mauritanum*) and *Solanum nigrum*.
- The groundcover was numerically dominated by the exotic grasses *Megathyrsus maximus* var. *maximus*, *Chloris gayana*, *Melinis repens*, Johnson's grass (*Sorghum halepense*) and vasey grass (*Paspalum urvillei*).
- Dominant exotic vines included *Ipomoea cairica*, *Macroptilium atropurpureum*, *Neonotonia wightii* and white passionflower (*Passiflora alba*). Sparse *Anredera cordifolia* was also recorded.
- Common exotic herbs and forbs included *Bidens pilosa*, lance-leaved rattlegod (*Crotalaria pallida*), gomphrena weed (*Gomphrena celosioides*), *Sida cordifolia*, purple-top (*Verbena bonariensis*), *Verbena litoralis*, *Sonchus oleraceus* and *Asclepias curassavica*.
- No aquatic macrophyte weed species were recorded and none are likely to occur given the brackish to saline conditions of the waterways.
- Except for recent clearing and associated weed removal for industrial development adjacent to the channel, no major differences in weed composition or density were observed at Port West between the current and previous monitoring surveys.

Results



Figure 3-3 Port West Drain 2018

3.3 Port Gate Drain

3.3.1 Site Description

Port Gate Drain located in the south-west section of the PBPL lies to the south of Boat Passage in the vicinity of Howard Smith Drive (refer to Figure 3-4). It collects stormwater run-off from the adjacent hardstand areas and drains into (and partially receives) the tidal waters in Boat Passage. The banks of the drain are constructed of concrete, gravel and/or compacted earth, which limits extensive vegetation growth. The tidal reaches of the drain support sparse fringing *Avicennia marina* seedlings, with a sparse groundcover of common reed (*Phragmites australis*) and widespread native sedges (*Schoenoplectus*, *Bolboschoenus*, *Fimbristylis*). Outside the tidal zone the banks support a sparse groundcover of exotic grasses and shrubs. RPS (2013) reports that Port Gate Drain undergoes regular maintenance (including mowing and spraying for exotic species) facilitated by the PBPL.

3.3.2 Weeds

The following observations were made in the 2018 survey:

- No DAWR-listed species were recorded within or directly adjacent to Port West Drain.
- Restricted Matters recorded included: *Baccharis halimifolia*, *Schinus terebinthifolius* and *Lantana camara*.
- Exotic species recorded were dominated by environmental weeds well established across Brisbane including: *Ipomoea cairica*, *Macroptilium atropurpureum*, *Chloris gayana*, *Megathyrsus maximus* var. *maximus*, *Bidens pilosa*, *Gomphocarpus physocarpus*, *Melinis repens*, *Solanum nigrum*, *Conyza bonariensis*, *Leucaena leucocephala*, *Araujia sericifera*, *Passiflora foetida* and *Macroptilium lathyroides*.
- The dominant woody weeds recorded included *Schinus terebinthifolius*, *Baccharis halimifolia* and *Lantana camara*. Other woody weeds included *Cinnamomum camphora*, *Celtis sinense*,

Results

Gomphocarpus physocarpus, *Leucaena leucocephala*, *Solanum chrysotrichum*, *Solanum nigrum* and *Psidium guajava*.

- The groundcover was numerically dominated by exotic grasses including *Megathyrsus maximus* var. *maximus*, *Chloris gayana*, *Melinis repens*, *Sorghum halepense*, *Cenchrus echinatus* and *Paspalum* spp. Other introduced grasses included *Arundo donax*, *Cynodon dactylon* and *Hyparrhenia rufa*.
- Dominant exotic vines included *Ipomoea cairica*, *Macroptilium atropurpureum*, *Vigna* sp., *Passiflora* spp. and *Araujia sericifera*.
- Common exotic herbs and forbs included *Bidens pilosa*, *Capsella* sp., *Emilia sonchifolia*, *Conyza bonariensis*, *Commelina benghalensis*, *Gomphrena celosioides*, *Sida cordifolia*, Singapore daisy (*Sphagneticola trilobata*) and *Tridax procumbens*.
- No aquatic macrophyte weed species were recorded and none are likely to occur given the brackish to saline conditions of the waterways.
- No major differences in weed composition were observed between the current and previous monitoring surveys at Port Gate Drain.



Figure 3-4 Port Gate Drain 2018

3.4 T1-3, Car Precinct and Constructed Lakes

3.4.1 Site Description

The T1-3 and Car Precinct areas east of Port Drive at the Port of Brisbane store imported vehicles and as such are potential vectors for newly introduced weed species (refer to Figure 3-5).

The T1-3 survey area extends around the car parking hardstand and includes a constructed drain to the north. The survey area consists predominantly of regularly maintained lawn which receives stormwater overflow from the hardstand. Landscaped garden beds front Port Drive. The constructed drain in the northern section consists of a concrete channel designed to pipe stormwater runoff into the Brisbane River.

Results

The Car Precinct survey area lies to the south and west of the hardstand vehicle storage area. The survey area consists predominantly of regularly maintained lawn which receives stormwater overflow from the hardstand.

The T1-3 and Car Precinct sites are separated by the Queensland Rail freight line and a cleared and regularly mowed easement. Constructed lakes lie south of the Car Precinct.

T1-3, Car Precinct, railway easement and lands surrounding the lakes undergo regular maintenance including mowing, and RPS (2013) reported that these areas were regularly sprayed for weeds.

3.4.2 Weeds

The following observations were made in the 2018 survey:

- No DAWR-listed species were recorded within or directly adjacent to the survey area.
- Restricted Plants recorded included *Baccharis halimifolia*, fireweed (*Senecio madagascariensis*), common ragweed (*Ambrosia artemisiifolia*), *Schinus terebinthifolius* and *Lantana camara*.
- Exotic species recorded were dominated by environmental weeds well established across Brisbane including: *Senecio madagascariensis*, *Schinus terebinthifolius*, *Lantana camara*, *Ipomoea cairica*, *Ricinus communis*, *Macroptilium atropurpureum*, *Cenchrus echinatus*, *Megathyrsus maximus* var. *maximus*, *Solanum nigrum*.
- Dominant woody weeds included *Schinus terebinthifolius*, *Baccharis halimifolia*, *Ricinus communis*, *Senecio madagascariensis*, *Lantana camara* and *Solanum nigrum*.
- Dominant exotic vines recorded included *Ipomoea cairica*, *Ipomoea alba* and *Macroptilium atropurpureum*.
- The sparse groundcover was regularly mowed. The most widespread exotic groundcovers were *Megathyrsus maximus* var. *maximus*, *Cenchrus echinatus*, *Melinis repens*, *Chloris gayana*, *Digitaria ciliaris*, *Cynodon dactylon*, *Paspalum urvillei*, *Setaria sphacelata* and *Setaria pumila*.
- Common exotic herbs and forbs included *Ageratum houstonianum*, *Cyperus eragrostis*, *Bidens pilosa*, *Crassocephalum crepidioides*, *Eclipta prostrata*, *Phytolacca octandra*, *Capsella bursapastoris*, *Gomphrena celosioides*, *Lantana montevidensis*, *Conyza bonariensis*, *Hypochaeris radicata*, *Oenothera drummondii*, *Tridax procumbens*, *Verbena bonariensis*, *Portulaca oleracea* and *Malvastrum coromandelianum*.
- Exotic aquatic macrophytes recorded in the shallow waters on the lakes edge included umbrella sedge (*Cyperus involucratus*) and *Ludwigia peploides*.
- No significant differences in weed composition between the current and previous monitoring surveys were observed at these sites.



Figure 3-5 T1-3, Car Precinct and Constructed Lake

3.5 The Bird Hide

The bird hide adjacent to Lucinda Drive is located on intertidal flats supporting a mix of mangroves, saltmarsh and mudflats which provide important habitat for a mix of migratory waders (refer to Figure 3-6). These intertidal conditions prohibit the establishment of most introduced weed taxa with the exception of minor patches of *Baccharis halimifolia* on elevated sites. As described above, whilst these high value intertidal habitats may be vulnerable to *Rhizophora mangle* invasion, this risk is considered low.

The filled lands above HAT support terrestrial grasslands which are regularly mown and support a range of grass and herbaceous environmental weeds widespread throughout the region. This includes *Macroptilium atropurpureum*, *Chloris gayana*, *Megathyrsus maximus* var. *maximus*, *Cenchrus echinatus*, *Stylosanthes scabra*, *Tridax procumbens*, *Gomphrena celosioides*, *Cynodon dactylon*, *Conyza bonariensis*, *Ipomoea cairica*, *Crassocephalum crepidioides*, *Setaria sphacelata*,

Results

Sonchus oleraceus, *Bidens pilosa*, *Portulaca oleracea*, *Ageratum houstonianum*, *Oenothera drummondii*, *Cakile edentula*, *Ipomoea cairica* and *Macroptilium lathyroides*. Sparse woody weeds include *Baccharis halimifolia* and *Lantana camara*.

Whilst the risk of weed invasion and associated habitat degradation within habitats of environmental value in the northern sector of the Port is considered low, with the exception of *Phragmites australis* spread into samphire and saltmarsh communities, site monitoring is recommended to protect the environmental values of these wetlands.



Figure 3-6 Bird Hide 2018

3.6 Port Drive Wetlands

Port lands south of Boat Passage and adjacent to Port Drive support a mix of freshwater and intertidal wetlands of high ecological value and are prone to varying levels of weed invasion (refer to Figure 3-8).

3.6.1 Intertidal Wetlands

An extensive mosaic of mangroves and saltmarsh occurs east of Port Drive. Whilst the mangroves are not prone to weeds due to regular tidal inundation, the saltmarsh-mangrove ecotone supported localised patches of *Schinus terebinthifolius* on higher ground with a groundcover of *Chloris gayana* and *Megathyrsus maximus* var. *maximus*.

As identified in previous monitoring surveys, the landward edge of the saltmarsh is regularly slashed for ground maintenance which is removing succulent saltmarsh species, altering local topography and promoting the spread of exotic grasses, particularly *Chloris gayana*, into saltmarsh habitat (refer to Figure 3-7). *Phragmites australis* also forms sparse to localised patches at the upper tidal limit of the saltmarsh. Whilst *Phragmites* naturally occurs at the freshwater-saline interface, it can spread rapidly to form extensive stands in saltmarsh and can alter the hydrology, sedimentation, ecology and function of the saltmarsh community, with flow on effects to adjacent intertidal habitat.

Results

All weed species recorded at this site are widespread across PBPL lands and are well-established in the Brisbane region and throughout coastal south-east Queensland. No DAWR-listed species were recorded within or directly adjacent to this survey area. However, the weed invasions at the mangrove-saltmarsh interface and at the upper saltmarsh limits are reducing the overall condition of these communities and providing local sources for weed spread. The regular slashing also reduces the condition of coastal saltmarsh and promotes habitat conditions more suitable for weed invasion.

3.6.2 Filled Intertidal Land

Approximately 1.0 ha of previous intertidal and low-lying land east of Port Drive has been partially filled and excluded from tidal inundation. The altered hydrology and topography has favoured regrowth dominated by terrestrial weed species including *Megathyrsus maximus* var. *maximus*, *Cenchrus echinatus*, *Melinis repens*, *Chloris gayana*, *Digitaria ciliaris*, *Paspalum* spp., *Pennisetum purpureum* and *Cortaderia selloana*. Patches of *Phragmites australis* also occur in lower lying depressions influenced by freshwater inundation. Whilst this is a native species, *Phragmites australis* can invade disturbed saltmarsh which has reduced tidal inundation and salinity and was observed invading some intact saltmarsh fringes across this site.

The exotic shrub *Schinus terebinthifolius* occurs as scattered individuals to dense thickets in this area depending on the level of inundation. Other exotic shrubs recorded in this filled and disturbed land include *Stylosanthes scabra*, *Opuntia stricta*, and *Solanum chrysotrichum*. Sparse exotic vines recorded included *Ipomoea cairica* and *Macroptilium atropurpureum*.

All weed species recorded at this disturbed site are widespread across PBPL lands and are well-established in the Brisbane region and throughout coastal south-east Queensland. No DAWR-listed species were recorded within or directly adjacent to this survey area. These weed-dominated habitats are likely to provide limited value for local fauna and estuarine species associated with the adjacent intertidal complex. In addition, these degraded habitats are replacing native communities, including coastal saltmarsh.

To enhance saltmarsh and reduce weed invasion in this area the ground level could be re-contoured to reinstate tidal inundation within the range suitable for saltmarsh establishment. Some level of variation in topography would need to be maintained during re-profiling to increase habitat diversity and to ensure excessive ponding does not occur. As the proposed works will disturb the ground profile, PASS assessments and mitigation measures would be required prior to any on-ground works. Natural recruitment is the preferred and most cost-effective method of revegetating re-profiled intertidal sites. However, active planting could be carried out to supplement natural regeneration and to fast-track the recovery process, if natural colonisation is slow.

2011:



2013:



2016:



2017:



Figure 3-7 Disturbed Coastal Saltmarsh Port Drive

3.6.3 Freshwater Wetlands

Freshwater wetlands adjacent to Port Drive support remnant *Melaleuca quinquenervia* dominated communities which are poorly represented on Port lands. Given their proximity to the rail and road corridors these wetlands are susceptible to weed invasion. However, these communities could not be surveyed in 2017 or 2018 due to construction of the Port Drive upgrade.



Figure 3-8 Port Drive Wetlands 2018

3.7 Port West Estuarine Wetlands

Port West supports an extensive mosaic of mangroves and saltmarsh directly connected to the Brisbane River (refer to Figure 3-9). Similar to other sites at the Port, the saltmarsh-mangrove ecotone and upper tidal limits, which lie adjacent to industrial land uses, are susceptible to disturbance and weed invasion.

The western landward fringe of closed mangrove forest comprises evenly aged *Avicennia marina* over a dense native saltmarsh shrublayer comprising *Sporobolus virginicus* and *Sesuvium portulacastrum*. Very sparse weed invasion was recorded under this dense mangrove canopy and comprised mainly isolated *Asparagus aethiopicus*, *Opuntia stricta*, *Schinus terebinthifolius*, *Ricinus communis*, *Leucaena leucocephala* and *Senna pendula* var. *glabrata* on elevated sites.

Locally elevated sites within saltmarsh landward of the mangrove fringe supported sparse *Schinus terebinthifolius* over a groundcover of *Megathyrsus maximus* var. *maximus* and *Chloris gayana*. More landward sites also supported *Senna pendula* var. *glabrata*, *Baccharis halimifolia*, *Ricinus communis*, *Opuntia stricta* and *Asparagus* spp. Some *Phragmites* invasion was also occurring at the landward edge of disturbed saltmarsh subject to freshwater ponding.

All weed species recorded within the saltmarsh-mangrove interface at Port West are widespread across PBPL lands and are well-established in the Brisbane region and throughout coastal south-east Queensland. No DAWR-listed species were recorded within or directly adjacent to this survey area.



Figure 3-9 Port West Estuarine Habitat 2018

3.8 Fort Lytton Estuarine Wetlands

In 2015, BMT WBM provided an assessment of a potential habitat improvement area at Fort Lytton on the eastern banks of the Brisbane River (refer to Figure 3-10). That assessment identified approximately 0.35ha of previously intertidal land at the ecotone between the mangroves and saltmarsh that had been filled and promoted weed invasion.

Re-survey of the site in 2017 and 2018 indicated that *Leucaena leucocephala* is becoming more widespread across this site. Other woody weeds included *Schinus terebinthifolius*, *Celtis sinense*, *Baccharis halimifolia*, *Phoenix* sp., *Lantana camara*, *Phytolacca octandra*, *Solanum nigrum*, *Gomphocarpus physocarpus*, *Opuntia stricta*, *Ricinus communis* and *Solanum mauritianum*. Dominant exotic groundcovers included *Chloris gayana*, *Megathyrsus maximus* var. *maximus* and *Sorghum halepense*. Other weeds recorded included *Anredera cordifolia*, *Ageratum houstonianum*, *Passiflora foetida*, *Conyza bonariensis*, *Capsella bursa-pastoris*, *Bidens pilosa*, *Cynodon dactylon*, *Gomphrena celosioides*, *Ipomoea cairica*, *Paspalum urvillei*, *Plantago lanceolata* and *Sida* spp. The placement of fill has also reduced tidal flow to approximately 0.06ha of saltmarsh which has promoted *Phragmites* invasion.

All weed species recorded at this site are widespread across PBPL lands and are well-established in the Brisbane region and throughout coastal south-east Queensland. No DAWR-listed species were recorded within or directly adjacent to this survey area.



Figure 3-10 Fort Lytton 2018

4 Discussion

In terms of weed composition and distribution, the survey sites have remained relatively stable over the monitoring period (2001-2018). With the exception of isolated *Hyparrhenia rufa* (a widespread grass weed in south-east Qld), no new weed species, or significant outbreaks of weed invasion, were recorded in the current survey.

Although all weeds recorded adjacent to Port Drive are widespread throughout south-east Queensland and no newly introduced or high risk species have been identified at this location, the road and rail corridors are potential vectors for newly imported species arriving at the Port and are important monitoring sites.

Several sites of saltmarsh disturbance have been recorded across PBPL lands at Port Drive, Port West and Fort Lytton. All sites have been subject to disturbance which has altered site hydrology and promoted weed invasion including the spread of *Phragmites australis*. The exotic shrubs *Schinus terebinthifolius* and *Leucaena leucocephala* also occur on pockets of higher ground in the saltmarsh-mangrove ecotone and are generally associated with a groundcover of exotic grasses particularly *Megathyrsus maximus* var. *maximus* and *Chloris gayana*. At the Port Drive and Fort Lytton sites the landward edge of the saltmarsh ecotone is also regularly slashed for ground maintenance which is removing succulent saltmarsh species, altering local topography and promoting the spread of exotic grasses into saltmarsh habitat.

It is recommended that woody weeds at the upper tidal limit are removed to reduce the source of local seed propagules and the local terrain is re-profiled to the range of saltmarsh and mangrove habitat. Regular slashing for site maintenance should be restricted to the area of terrestrial grasses only, with a minimum 10 m buffer provided to saltmarsh habitat. In addition to the threat of site maintenance, coastal saltmarsh on Port lands is vulnerable to sea level rise and associated mangrove expansion.

Figure 4-1 indicates the extent of mangrove encroachment in to coastal saltmarsh at Port Drive between 2009 and 2018. There has been no significant increase in coastal saltmarsh in this area over the same period.

Re-profiling highly disturbed areas of exotic grassland adjacent to tidal lands could assist saltmarsh resilience to sea level rise and may also promote *Casuarina glauca* recruitment. *C. glauca* communities in south-east Qld have recently been listed as Endangered under the *EPBC Act*. Some level of variation in topography should be maintained during re-profiling to increase habitat diversity and to ensure excessive ponding does not occur. PASS assessments and mitigation measures will also be required prior to any on-ground works. Natural recruitment is the preferred and most cost-effective method of revegetating re-profiled intertidal sites. However, active planting could be carried out to supplement natural regeneration and to fast-track the recovery process.

It is recommended that future annual weed surveys are carried out at the same sites and in accordance with the rapid biological assessment methodologies described in this report.

Discussion



Figure 4-1 Mangrove Encroachment of Coastal Saltmarsh Port Drive 2009-2018

5 Conclusion

All weed species recorded on PBPL lands are well-established in the Brisbane region and throughout coastal south-east Queensland. No targeted DAWR-listed species were recorded within or directly adjacent to the survey areas.

Weed invasion on PBPL lands is associated with site disturbance, or in habitats directly adjacent to disturbance, on terrestrial lands and at the limit of HAT. There is no weed invasion within habitats subject to regular tidal inundation. The marine weed species *Rhizophora mangle* has been recorded in Australia, but has not been observed in south-east Queensland, and is considered a low risk species. Despite the low risk, regular weed monitoring should be maintained to ensure this species, and potential high risk taxa, do not establish in Port habitats.

Survey sites have remained relatively stable over the weed monitoring period (2001-2018). No new weed species or sites of significant weed incursion and spread were detected. Several sites of saltmarsh disturbance were recorded across PBPL lands where altered site hydrology has promoted weed invasion, particularly *Phragmites australis*, *Schinus terebinthifolius* and *Leucaena leucocephala*. At some sites the landward edge of the saltmarsh ecotone is also regularly slashed for ground maintenance which is promoting the spread of exotic grasses. Ground works and changing land maintenance practices at these sites could reinstate habitat conditions more suitable for native saltmarsh species, including *Casuarina glauca*. In addition to the threat of site maintenance, coastal saltmarsh in the Port Drive area may be vulnerable to sea level rise and associated mangrove expansion. Re-profiling the area of exotic grassland in this area could improve saltmarsh resilience to sea level rise.

The weed species recorded on PBPL lands are widespread in Brisbane and the south-east Queensland bioregion, and for that reason will be difficult to control and are unlikely to be eradicated from PBPL lands. Despite this, it is recommended that PBPL continue weed control efforts to fulfil landholder obligations under the *Biosecurity Act*. This includes monitoring the Port Drive corridor and sites of high value habitat, particularly saltmarsh which is vulnerable to disturbance associated with land use. Ongoing management will also reduce potential future costs associated with delaying weed control, particularly the removal of mature woody shrubs and dense vine infestations. In addition, active weed control and habitat restoration in disturbed saltmarsh would improve the habitat condition of this locally restricted and nationally recognised habitat of high ecological value.

6 References

- BCC (2013). Brisbane Invasive Species Management Plan 2013-17. Prepared by the Brisbane City Council, February 2013.
- BMT WBM (2014). Port of Brisbane Annual Weed Survey – 2014. Report prepared for PBPL.
- BMT WBM (2015). Port of Brisbane Annual Weed Survey – 2015. Report prepared for PBPL.
- BMT WBM (2016). Port of Brisbane Annual Weed Survey – 2016. Report prepared for PBPL.
- BMT WBM (2017). Port of Brisbane Annual Weed Survey – 2017. Report prepared for PBPL.
- Brisbane City Council (2014). Weed Classification. <http://weeds.brisbane.qld.gov.au/weed-classification>, accessed August, 2014.
- Queensland Herbarium (2016). Intertidal Communities of Moreton Bay. Change in Extent 1955-1997-2012 Mangrove and Associated Communities of Moreton Bay Queensland Australia.
- RPS Australia East Pty Ltd (2013). Annual Plant Assessment Port of Brisbane. Report Prepared for the Port of Brisbane, July 2013.

Appendix A Dominant Weed Species Recorded at PBPL

Species	Common Name	Lifeform
<i>Ageratum houstonianum</i>	Blue billygoat weed	forb
<i>Alternanthera denticulata</i>	Lesser joyweed	forb
<i>Bidens pilosa</i>	Cobbler's pegs	forb
<i>Commelina benghalensis</i>	Hairy commelina	forb
<i>Conyza bonariensis</i>	Flaxleaf fleabane	forb
<i>Crassocephalum crepidioides</i>	Thickhead	forb
<i>Einadia hastata</i>	Berry saltbush	forb
<i>Emilia sonchifolia</i>	Emilia	forb
<i>Impatiens</i> sp.	Balsam	forb
<i>Lantana montevidensis</i>	Creeping lantana	forb
<i>Ludwigia longifolia</i>	Long-leaved willow primrose	forb
<i>Onopordum acanthium</i>	Scotch thistle	forb
<i>Plantago lanceolata</i>	Ribwort	forb
<i>Senecio madagascariensis</i>	Fireweed	forb
<i>Sida rhombifolia</i>	Common sida	forb
<i>Sonchus oleraceus</i>	Common sowthistle	forb
<i>Sphagneticola trilobata</i>	Singapore daisy	forb
<i>Stachytarpheta jamaicensis</i>	Light blue snakeweed	forb
<i>Tagetes minuta</i>	Stinking Roger	forb
<i>Tridax procumbens</i>	Tridax daisy	forb
<i>Verbena bonariensis</i>	Purpletop	forb
<i>Datura</i> sp.	Thorn apple	forb
<i>Andropogon virginicus</i>	Whisky grass	grass
<i>Cenchrus echinatus</i>	Mossman River grass	grass
<i>Chloris gayana</i>	Rhode's grass	grass
<i>Cynodon dactylon</i>	Green couch	grass
<i>Digitaria ciliaris</i>	Summer grass	grass
<i>Hyparrhenia rufa</i>		grass
<i>Megathyrsus maximus</i> var. <i>maximus</i>	Green panic	grass
<i>Paspalum dilatatum</i>	Paspalum	grass
<i>Paspalum distichum</i>	water couch	grass
<i>Phragmites australis</i>	Common reed	grass
<i>Setaria sphacelata</i>	South African pigeon grass	grass

Dominant Weed Species Recorded at PBPL

Species	Common Name	Lifeform
<i>Sorghum halepense</i>	Johnson grass	grass
<i>Sporobolus fertilis</i>	Giant Parramatta Grass	grass
<i>Sporobolus virginicus</i>	Sand couch	grass
<i>Cyperus exaltatus</i>	Tall flatsedge	sedge
<i>Cyperus involucratus</i>	Umbrella sedge	sedge
<i>Baccharis halimifolia</i>	Groundsel	shrub
<i>Crotalaria pallida</i>	Rattlepod	shrub
<i>Lantana camara</i>	Lantana	shrub
<i>Opuntia stricta</i>	Prickly Pear	shrub
<i>Phytolacca octandra</i>	Inkweed	shrub
<i>Rivina humilis</i>	Coral berry	shrub
<i>Solanum chrysotrichum</i>	Giant Devil's fig	shrub
<i>Solanum nigrum</i>	Blackberry nightshade	shrub
<i>Celtis sinensis</i>	Chinese Elm	tree
<i>Cinnamomum camphora</i>	Camphor laurel	tree
<i>Ficus elastica</i>	Rubber fig	tree
<i>Leucaena leucocephala</i>	Leucaena	tree
<i>Ricinus communis</i>	Castor oil tree	tree
<i>Schinus terebinthifolius</i>	Broad-leaf Pepper Tree	tree
<i>Senna pendula</i> var. <i>glabrata</i>	Easter cassia	tree
<i>Solanum mauritianum</i>	Wild tobacco	tree
<i>Anredera cordifolia</i>	Madeira vine	vine
<i>Asparagus aethiopicus</i> cv. <i>Sprengeri</i>	Basket asparagus fern	vine
<i>Cardiospermum grandiflora</i>	Balloon vine	vine
<i>Ipomoea cairica</i>	Mile-a-minute	vine
<i>Macroptilium atropurpureum</i>	Siratro	vine
<i>Neonotonia wightii</i>	Glycine	vine
<i>Passiflora foetida</i>	Stinking passionflower	vine
<i>Passiflora suberosa</i>	Corky passionflower	vine
<i>Solanum seaforthianum</i>	Brazilian nightshade	vine

BMT has a proven record in addressing today's engineering and environmental issues.

Our dedication to developing innovative approaches and solutions enhances our ability to meet our client's most challenging needs.



Brisbane

Level 8, 200 Creek Street
Brisbane Queensland 4000
PO Box 203 Spring Hill QLD 4004
Tel +61 7 3831 6744
Fax +61 7 3832 3627
Email brisbane@bmtglobal.com

Melbourne

Level 5, 99 King Street
Melbourne Victoria 3000
Tel +61 3 8620 6100
Fax +61 3 8620 6105
Email melbourne@bmtglobal.com

Newcastle

126 Belford Street
Broadmeadow New South Wales 2292
PO Box 266 Broadmeadow
New South Wales 2292
Tel +61 2 4940 8882
Fax +61 2 4940 8887
Email newcastle@bmtglobal.com

Adelaide

5 Hackney Road
Hackney Adelaide South Australia 5069
Tel +61 8 8614 3400
Email info@bmttdt.com.au

Northern Rivers

Suite 5
20 Byron Street
Bangalow New South Wales 2479
Tel +61 2 6687 0466
Fax +61 2 6687 0422
Email northernrivers@bmtglobal.com

Sydney

Suite G2, 13-15 Smail Street
Ultimo Sydney New South Wales 2007
Tel +61 2 8960 7755
Fax +61 2 8960 7745
Email sydney@bmtglobal.com

Perth

Level 4
20 Parkland Road
Osborne Park WA 6017
PO Box 2305 Churchlands WA 6918
Tel +61 8 6163 4900
Email perth@bmtglobal.com

London

1st Floor, International House
St Katharine's Way
London
E1W 1UN
Tel +44 (0) 20 8090 1566
Email london@bmtglobal.com

Aberdeen

Broadfold House
Broadfold Road, Bridge of Don
Aberdeen
AB23 8EE
UK
Tel: +44 (0) 1224 414 200
Fax: +44 (0) 1224 414 250
Email enquiries@bmtcordah.com

Asia Pacific

Indonesia Office
Perkantoran Hijau Arkadia
Tower C, P Floor
Jl: T.B. Simatupang Kav.88
Jakarta, 12520
Indonesia
Tel: +62 21 782 7639
Fax: +62 21 782 7636
Email asiapacific@bmtglobal.com

Alexandria

4401 Ford Avenue, Suite 1000
Alexandria
VA 22302
USA
Tel: +1 703 920 7070
Fax: +1 703 920 7177
Email inquiries@dandp.com